Water footprint analysis (hydrologic and economic) of the Guadiana river basin1

M.M. Aldaya $^{\rm a}$ and M.R. Llamas $^{\rm b}$

E-mail: m.m.aldaya@ctw.utwente.nl

E-mail: mrllamas@geo.ucm.es

1 Admitted as an accompanying report to the U.N. World Water Development Report No. 3 to be presented during the 5th World Water Forum (Istanbul, March, 2009)

^a Twente Water Centre. University of Twente. The Netherlands

^b Department of Geodynamics. Complutense University of Madrid, Spain

Table of contents

Sui	mmary	3
1.	Introduction	5
2.	Study area	8
3.	Methodology	9
4.	Data sources and limitations	12
5.	Results	14
	5.1 Guadiana water footprint	15
	A. Crop area	15
	B. Water consumption	16
	C. Virtual water content (m ³ /ton)	17
	D. Agricultural economic productivity (€ha)	19
	E. Economic blue water productivity (€m³)	20
	F. Agricultural trade	22
	5.2 Review of crop water consumption estimates	24
6.	Conclusions	25
	Acknowledgements	28
	Symbols	29
	Glossary	30
	Bibliography	33
	Appendices	41
	I. Crop area, production and yield	41
	II. Crop water consumption and virtual water content	41
	III. Economic value and economic water productivity	41

Summary

In most arid and semiarid countries, water resources management is an issue as important as controversial. Today most water resources experts admit that water conflicts are not caused by the physical water scarcity but they are mainly due to poor water management or governance. The virtual water concept, defined as the volume of water used in the production of a commodity, good or service, together with the water footprint (water volume used to produce the goods and services consumed by a person or community), link a large range of sectors and issues, providing an appropriate framework to find potential solutions and contribute to a better management of water resources, particularly in arid or semi-arid countries.

As the most arid country in the European Union, water use and management in Spain is a hot political and social topic. The aim of this study is to analyse the virtual water and water footprint, both from a hydrological and economic perspective, in the semiarid Guadiana basin. The transboundary Guadiana river basin located in south-central Spain and Portugal drains an area of 66,800 km², of which 17% lies in Portugal. The present analysis is carried out in the Spanish side of the basin which has been divided in Upper, Middle, Lower Guadiana basin and TOP domain. The TOP domain is a group of three small river basins located near the Guadiana River mouth. In these regions the main green and blue water consuming sector is irrigation, with about 95% of total water consumption. Within this sector, high virtual-water low-economic value crops are widespread in the studied Upper and Middle Guadiana regions, particularly cereals with low blue water economic productivity. In particular, the Upper Guadiana basin is among the most significant in Spain in terms of conflicts between agriculture, with almost no food (virtual water) import, and

the conservation of rivers and groundwater-dependent wetlands. On the other hand, in the Lower Guadiana basin and TOP domain, growing vegetables and crops under plastic greenhouses, the blue water economic productivity values are much higher, using jointly surface and groundwater resources. The amount of crops and the employment generated in the whole Guadiana basin is already producing 'more crops and jobs per drop'. The aim now is to move towards the policy 'more cash and nature per drop', especially in the Upper and Middle Guadiana basin.

1. Introduction

In most arid and semiarid countries, water resource management is an issue as important as controversial. Today most water resources experts admit that water conflicts are not caused by the physical water scarcity but they are mainly due to poor water management. The virtual water and water footprint analysis, linking a large range of sectors and issues, provides an appropriate framework to find potential solutions and contribute to a better management of water resources, particularly in water scarce countries.

The water footprint (WF) is a consumption-based indicator of water use defined as the total volume of water that is used to produce the goods and services consumed by an individual or community (Hoekstra and Chapagain, 2008). Closely linked to the concept of water footprint is the virtual water. The virtual water content of a product (a commodity, good or service) refers to the volume of water used in its production (Allan, 1997; 1999; Hoekstra, 2003). Building on this concept, virtual water 'trade' represents the amount of water embedded in traded products (Hoekstra and Hung, 2002). A nation can preserve its domestic water resources by importing water intensive products instead of producing them domestically (ibid.). These 'water savings' can be used to produce alternative, higher-value agricultural crops, to support environmental services, or to serve growing domestic needs. Thus, virtual water 'import' is increasingly perceived as an alternative source of water for some water-stressed nations and is starting to change the current concepts of water and food security.

Furthermore, the virtual water and water footprint analysis makes explicit how much water is needed to produce different goods and services. In semi-arid and arid areas, knowing the virtual water value of a good or service can be useful towards determining how best to use the scarce water available. In this sense, it is important to establish whether

the water used proceeds from rainwater evaporated during the production process (green water) or surface water and/or groundwater evaporated as a result of the production of the product (blue water) (Chapagain et al., 2006; Falkenmark, 2003). Traditionally, emphasis has been given to the concept of blue water through the 'miracle' of irrigation systems. However, an increasing number of authors highlight the importance of green water (Allan, 2006; Comprehensive Assessment of Water Management in Agriculture, 2007; Falkenmark and Rockström, 2004; Rockström, 2001). Virtual water and water footprint assessment could thus inform production and trade decisions, promoting the production of goods most suited to local environmental conditions and the development and adoption of water efficient technology. Adopting this approach, however, requires a good understanding of the impacts of such policies on socio-cultural, economic and environmental conditions. Besides, water is not the only factor of production and other factors, such as energy, may come to play an increasingly important role in determining water resources allocation and use.

The present study deals with the economic and hydrological analysis of the virtual water and water footprint of the Guadiana river basin, considering both green and blue (ground and surface) water of the different economic sectors. This could facilitate a more efficient allocation and use of water resources, providing simultaneously a transparent interdisciplinary framework for policy formulation. The Guadiana river basin is shared by Spain and Portugal, but this report focuses on the Spanish area of the river basin. The analysis of the Portuguese area (less than 20% of the total area of the basin) will be carried out by the Portuguese INAG (National Water Authority). It analyses the water footprint, virtual water and economic relevance of each economic sector in different rainfall years (evaluating an average - 2001, dry -2005, and humid year -1997). Special emphasis is given

to the agricultural sector, which consumes about 95% of total green and blue water resources. First of all, the whole Guadiana is evaluated, which has been divided in four sections: groundwater based Upper Guadiana basin, mainly surface water based Middle basin, both groundwater and surface water based Lower Guadiana basin and the former Lower Guadiana or Guadiana II (henceforth TOP domain)² comprising the Tinto, Odiel and Piedras river basins. At the end of the chapter virtual water 'trade' is evaluated. Finally, crop water consumption estimates are assessed against the results obtained by other national and international studies. A glossary with key terms is also included at the end of the study, a more extensive version can be found in (Aldaya and Llamas, in press). It concludes that a better knowledge of the water footprint and virtual water 'trade' in the semiarid Guadiana basin provides a transparent and multidisciplinary framework for informing and optimising water policy decisions, contributing at the same time to the implementation of the EU Water Framework Directive (2000/60/EC). As a whole, the Guadiana river basin has already achieved a good degree of the paradigm 'more crops and jobs per drop' but it is still far from achieving 'more cash and nature per drop'. An exception for this is the case of the Lower Guadiana basin and TOP domain in Andalusia, where virtual water-extensive high economic value crops adapted to the Mediterranean climate are grown, essentially vegetables, fruits and olive oil. For the time being and almost in the entire world, water footprint analysis has focused on hydrological aspects. A significant innovation of this work is to emphasize the imperative challenge of considering economic and ecological aspects, with the aim of going towards the new paradigm 'more cash and nature per drop' (Aldaya et al., 2008). Finally, the water footprint analysis is

² Before 1 January 2006 the TOP domain was the competence of the Guadiana River Basin Authority, after this date it was transferred to the Government of Andalusia.

providing new data and perspectives that are enabling to get a more optimistic outlook of the frequently spread looming «water scarcity crisis». We expect that this new knowledge makes traditional water and food security concepts change, concepts that have hitherto prevailed in the minds of most policy makers.

2. Study area

The Guadiana basin has an area of about 67,000 km² (83% in Spain and 17% in Portugal). The climate is semiarid, with an average precipitation of about 450 mm/year and average annual temperature of 14-16 °C (CHG, 2008a; INAG, 2007).

For practical purposes, the basin has been divided in four areas (Figure 1): a) groundwater based Upper Guadiana basin (totally located in a part of the Castilla-La Mancha Autonomous region); b) mainly surface water based Middle Guadiana basin (comprising part of Extremadura but not the small fraction of Cordoba); c) the Lower Guadiana basin (including the part of the basin in Huelva); and d) TOP domain (comprising the Tinto, Odiel and Piedras river basins). The TOP domain was the competence of the Guadiana River Basin Authority before 1 January 2006, but its competence was then transferred to the Government of Andalusia (CHG, 2008a).

Figure 1. Guadiana river basin geographic and administrative domain from 1 January 2006 onwards (CHG, 2008a)

According to CHG (2008b) when referring to the Guadiana river basin on the whole ('Total Guadiana' in the present document), it includes the Upper, Middle and Lower basins including the small fraction of Cordoba.

Figure 2. Western Mancha aquifer location within the Upper Guadiana Basin. Modified from CHG (2008b).

The Upper Guadiana basin, located in Castilla-La Mancha, is one of the driest river basins in Spain (Hernández-Mora et al., 2003). In this part, UNESCO recognized the collective ecological importance of 25,000 ha of wetlands in 1980, when it designated the 'Mancha Húmeda' Biosphere Reserve. In a largely arid region, these wetlands provided crucial nesting and feeding grounds for European migrating bird populations and were home to rare animal and plant species. The Tablas de Daimiel National Park (2,000 ha), a Ramsar Site, stands out for its significance as a symbol for the Spanish conservation movement. Today, however, this wetland that used to receive the natural discharge from the Western Mancha aquifer (Figure 2), survive artificially, in a kind of 'ecological coma', thanks to the water transfers that come from the Tagus-Segura Aqueduct starting in 1988 (Hernández-Mora et al., 2003) and to the artificial pumpage of groundwater to maintain flooded about the 5% of the 2,000 hectares of wetlands in the undisturbed National Park. More recently, some NGOs are claiming that 'La Mancha Humeda, Biophere Reserve' should not be considered any more by UNESCO as a World Biosphere Reserve. On the other hand, in order to recover these ecosystems, the Spanish Government, at the proposal of the Ministry of the Environment, approved a Special Plan for the Upper Guadiana (Plan Especial del Alto Guadiana -PEAG) on 11 January 2008 (CHG, 2008c). The formal approval of this Plan includes a budget of 5,500 million euros to be spent during the next 20 years.

3. Methodology

The present study estimates the virtual water and water footprint of the Guadiana river basin considering the green and blue water components for the most representative crops and the blue water component for livestock, industrial products and domestic (urban) water use. Within the blue water component, the volumes of surface and groundwater consumption are differentiated. In parallel with these analyses, economic data are studied.

This is done for each section of the river basin (Upper, Middle, Lower Guadiana and TOP domain) at different time scales (for an average -2001, dry -2005, and humid year -1997).

The virtual water and water footprint are calculated using the methodology developed by Hoekstra and Hung (2002; 2005) and Chapagain and Hoekstra (2003; 2004). For its emphasis on green and blue water, the present research follows recent works of Chapagain et al. (2006) Hoekstra and Chapagain (2008) and Chapagain and Orr (2008).

The method followed in this work is described in more detail in a more extensive report of the Guadiana river basin (Aldaya and Llamas, in press).

Virtual water content

The virtual water content of a product is the volume of freshwater used to produce the product, which depends on the water use in the various steps of the production chain. The virtual water content of primary crops (m³/ton) (i.e. crops in the form as they come directly from the land without having undergone any processing) has been calculated at a provincial level as the ratio of the volume of water used during the entire period of crop growth (crop water use, m³/year) to the corresponding crop yield (ton/ha) in the producing region. The crop water requirement of a certain crop under particular climatic circumstances was estimated with the CROPWAT model (Allen et al., 1998; FAO, 2003) using climate data at a provincial level. The volume of water used to grow crops in the field has two components: effective rainfall (green water) and irrigation water (blue water).

One particularity of the methodology used in this detailed study is that rainfed and irrigated agriculture were differentiated in the green and blue virtual water component calculations. That is, the green component in the virtual water content of a primary crop $(V_g, \, \text{m}^3/\text{ton})$ is calculated as the green crop water use (m^3/ha) divided by the crop yield $(Y_g, \, \text{ton/ha})$ for both rainfed $(Y_g, \, \text{m}^3/\text{ton})$ and irrigated production (Y_{irr}) . In parallel, the blue water

component $(V_b, \text{ m}^3/\text{ton})$ is calculated as blue crop water use divided by crop yield in irrigated production (Y_{irr}) .

Water footprint

In line with Chapagain and Hoekstra (2004), the water footprint of a country is equal to the total volume of water used, directly or indirectly, to produce the goods and services consumed by the inhabitants of the country. A national water footprint has two components, the internal and the external water footprint. First, the internal water footprint is defined as the volume of water used from domestic water resources to produce the goods and services consumed by the inhabitants of the region (Hoekstra and Chapagain, 2008). It is the sum of the total water volume used from the domestic water resources in the national economy minus the volume of virtual water export to other countries insofar related to export of domestically produced products. Second, the external water footprint is the volume of water used in other regions to produce goods and services imported and consumed by the inhabitants of that region. The present study calculates the water footprint per sub-basin related to production. Trade data at a provincial level are presented separately.

4. Data sources and limitations

In order to carry out this report, a number of simplifications have been assumed. First of all, the virtual water content values obtained with the CROPWAT model should be considered as a first approximation to reality. The main gaps in this approach are: a) the lack of data on the soils characteristics and their storage capacity for the effective rain; b) the amount of irrigation water 'lost' from the surface reservoirs to the field; c) the amount of water necessary to abate the pollution; and d) the reduction in crop yield when the irrigation demand cannot be supplied. Second, the eight most representative crops in each area have been studied corresponding to about 80% of the total area (Appendix 1). Third, with the aim of analysing the impact of climate variability on the use of water resources three different rainfall years were chosen: a humid (1997), average (2001) and dry year (2005). The average rainfall in 2001 was about 355 mm in Castilla-La Mancha, 547 in Extremadura and and 510 mm in Andalucía. When available, data for these years were used. This was not possible, however, in every case as shown below in this chapter. Fourth, and following CHG (2008b) data, when estimating the urban water use, urban water supply and sanitation data have been taken into account. Fifth, concerning the industrial water use, since energy and building industry are not considered within the industrial sector, hydroelectric energy was not included (CHG, 2008b). Sixth, with regard to the livestock water consumption, the drinking water and water to clean its housing is considered, leaving out the water used to grow and process its fodder. This is important when comparing these data with other analyses of the livestock water footprint. Finally, data have been compiled from different sources.

Geographic and social data: Data related to human population and employment were taken from the Guadiana River Basin Authority (CHG, 2008b).

Climatic data: Average monthly rainfall and evapotranspiration data at provincial level, as an input for the CROPWAT model (FAO, 2003), were obtained from the National Institute of Meteorology (INM, 2007).

Agricultural data: Data related to area (total area, crop area both rainfed and irrigated, irrigated area by irrigation system) were taken from the Guadiana River Basin Authority (CHG, 2008b) and the Spanish Ministry of Agriculture, Fisheries and Food 1T sheets (MAPA, 1999; 2001b).

Data on average rainfed and irrigated crop yield (*Y*) (kg/ha) at provincial level were taken from the Agro-alimentary Statistics Yearbook of the Spanish Ministry of Agriculture, Fisheries and Food (MAPA, 2007).

With regard to the crop parameters, as input data to CROPWAT, the crop coefficients in different crop development stages (initial, middle and late stage) were taken from FAO (Allen et al., 1998; FAO, 2003). The length of each crop in each development stage was obtained from FAO (Allen et al., 1998; FAO, 2003) when the climate region was specified; otherwise it was obtained from the work of Chapagain and Hoekstra (2004). The crop calendar was taken from the Spanish Ministry of Agriculture, Fisheries and Food (MAPA, 2001a). These data are also given at provincial level.

Economic data: Data related to gross value added (GVA) were taken from the Guadiana River Basin Authority (CHG, 2008b). Gross Value Added is obtained by deducting intermediate consumption from final agricultural production. That is, the gross value added is equal to net output or benefit to the farmer that can be used for the remuneration of productive factors. Nevertheless, in this study we will focus on the final economic agricultural production (total €) as well.

Crop economic value (€ton) for the different years was obtained from the Spanish Ministry of Agriculture, Fisheries and Food (MAPA, 2007). We are aware, however, that prices may change significantly from one year to the other. These data are an average for the whole Spain. In the present report CAP subsidies were not included (CHG, 2008b).

Hydrologic data: Data related to water origin (surface and groundwater) by agricultural region were taken from the Guadiana River Basin Authority (CHG, 2008b), which is based on the 1999 Agrarian Census of the National Statistics Institute (INE, 2007).

Green and blue crop consumptive water use (CWU, m³/ha) data were estimated using the CROPWAT model (FAO, 2003) (see Methodology section). Data on blue water withdrawals (surface and ground water) were taken from the Guadiana River Basin Authority (2007). It is noteworthy that these withdrawals are not the same as the estimated water consumption or evapotranspirative demand.

Average global irrigation efficiency at provincial level was taken from the CHG (2008b). It depends on the type of irrigation technique used by the farmer. Localized or drip irrigation is the most efficient system with a 0.9 coefficient, followed by sprinkler irrigation with 0.7 and finally, surface flood irrigation with 0.5.

Trade data: Data related to international trade at a provincial level were taken from ICEX (2008).

5. Results

Since irrigated agriculture is the main blue water user in the Guadiana Basin (about 90% according to MIMAM, 2007), the present study mainly focuses on water use by this sector. First of all, as seen in the methodology chapter, the Guadiana river basin has been divided and analyzed in four areas (Upper, Middle, Lower Guadiana and TOP domain). Then, the

obtained green and blue crop water consumption values are compared with national and international studies.

5.1. Guadiana water footprint

When comparing the Guadiana basin Gross Value Added (GVA) with national figures for the different sectors, the agricultural sector represents a value of 8.4 % of the national total, having both agriculture and livestock similar shares. Agriculture of the TOP domain represents 1.6 % of the national GVA, representing the livestock just a small amount (0.3 %). Concerning the manufacture industrial sector GVA, both in the Guadiana basin and TOP domain, it is not relevant in comparison with the total national, representing 1.99 % and 0.45 % of the total national respectively. These figures show the relevance of agriculture in these areas in comparison with other Spanish regions where industry and tourism are more important.

A. Crop area

The Spanish Guadiana river basin crop area is 26,000 km², which is about 47% of the total area. As a whole, in the basin, 19% of the crop area is devoted to irrigated agriculture. This proportion is similar to the Spanish average which amounts to 22% (MIMAM, 2007).

As shown in figure 3, the area dedicated to each crop type varies in each Guadiana section in the year 2001 (average precipitation). When looking at the rainfed agriculture similar crops are grown in the different Guadiana sections, highlighting cereals, olive trees and vineyards. Concerning irrigated agriculture, in general, cereals, vineyards and olive trees dominate in the Upper and Middle Guadiana basins, whereas citrus trees and vegetables in the Lower Guadiana and TOP domain. After the Common Agricultural Policy reform (2003), however, vineyard and olive tree irrigated production has increased significantly in Spain (18% y 16% respectively) (MAPA, 2006). According to Garrido and Varela (2008)

this is notable in Castilla- La Mancha Autonomous Community. It is expected that significant changes in crop distribution will continue to occur in the near future due to different causes, such as the increase in cereal prices.

Figure 3. Percentage of areas of irrigated and rainfed crops in the Upper, Middle, Lower Guadiana and TOP domain (average-year 2001). Showing crops occupying over 1% of land. Source: CHG (2008b).

B. Water use and consumption: total and by the agricultural sector

Total Water Use

As in most arid and semiarid regions, in the Guadiana river basin the main green and blue water consuming sector is irrigation, with about 95% of total water consumption in the basin as a whole (Table 1). The following main blue water user is urban water supply with less than 5% of the water applied for irrigation. If we consider that most urban water returns to the system, it can be said that irrigation consumptive uses are more than 95% of all the uses. However, the security of this supply is extremely relevant from a political and economic point of view. Concerning the Andalusian part (Lower Guadiana and the so-called TOP domain), irrigation consumes a lower water proportion, of about 75-80%, which account for the increase of the urban water supply. The industrial sector, even if it is the smallest water user, represents the highest economic value (GVA). Agriculture is also a significant economic activity in the Guadiana river basin, being the most important share of the GVA after the industrial sector (Table 1). Thus, even if urban and industrial uses have an obvious economic and social relevance, agriculture, as the highest water consumer in the basin, is the key to water resources management in the area.

Concerning rainfed and irrigated farming in the whole basin excluding TOP domain, total rainfed area is more than five times the irrigated area $(2,100x10^3)$ and $(2,100x10^3)$ hectares

respectively) (Appendix 2). Rainfed systems consume about 55% of the total water consumed by the agricultural sector (Table 1) and use green water (i.e. rainfall) that has a lower opportunity cost compared to the blue water use (i.e. irrigation) (Chapagain et al., 2005). Even if significantly smaller in extension, irrigated agriculture produces more tonnes and euros than rainfed agriculture (Appendix 2A and 2C).

Table 1. Internal water footprint of the Guadiana basin (year 2001)

Agricultural water consumption

As shown in figure 4, when taking into account rainfed and irrigated water consumption, crop water requirements are somewhat higher in the humid year. As it might be expected, there are remarkable variations in the green and blue water proportions in years with different rainfall patterns, being the blue water consumption higher in dry years and lower in humid years. While logically the green water consumption shows the opposite pattern.

Figure 4. Theoretical green and blue agricultural water consumption (Mm³/year) in the Upper, Middle, Lower Guadiana and TOP domain a dry (2005), average (2001) and humid year (1997). Source: Own elaboration.

The blue water consumption in the Upper Guadiana basin is mainly based on its groundwater resources, whereas the Middle Guadiana basin uses its surface water resources, mainly coming from large surface water reservoirs (Figure 5). The Lower Guadiana basin and TOP domain combine both ground and surface water strategies.

Figure 5. Theoretical green and blue (surface and ground) agricultural water consumption (Mm³/year) in the Upper, Middle, Lower Guadiana and TOP domain a dry (2005), average (2001) and humid year (1997). The size of the circle is proportional to the volume of water.

Source: Own elaboration.

C. Virtual water content in irrigated lands (m³/ton)

The virtual water analysis establishes the amount of water required by specific crops and it differs considerably among crop and climate types. For instance, Spain has a comparative advantage over most of the other European countries in the production of Mediterranean crops (such as vegetables, citrus fruits, vineyards or olive oil). It is also important to determine whether the water used proceeds from blue (i.e. irrigation) or green water (i.e. rainfall), and whether the blue water is surface or ground water.

Figure 6 provides an overview of the virtual water content of irrigated crops (m³/ton) in the different sections of the Guadiana basin in the different rainfall years. As shown in this figure, it is noteworthy that, among the studied crops, industrial crops (such as sunflowers), grain legumes, grain cereals (1,000-1,300 m³/ton) and olive trees (about 1,000-1,500 m³/ton) show the highest virtual water contents in irrigated agriculture. In humid years, however, olive trees are mainly based on green water resources. As previously mentioned, until recently, olive trees (and vineyards) were typical rain-fed crops. However, in last years the irrigated area seems to be significantly increasing for both crops.

It is widely believed that maize and vegetables are water-wasteful since in terms of m³/ha these crops consume large amounts of water. Nevertheless, when looking at the virtual water content in m³/kg these crops consume less water than it is generally believed. In fact, among the studied crops vegetables (100-200 m³/ton) exhibit the smallest virtual water content figures, probably due to the high yields they have.

Finally, vineyards have intermediate virtual water contents, of about 300-600 m³/ton.

Despite the semiarid nature of the Guadiana basin, in the Upper and Middle Guadiana basin irrigated grain cereal production is widespread in the year 2001. Aside from cereals, vineyards and olive trees are the most widespread crop in the basin during the year 2001. Two reasons may explain this trend. First, vineyards are significantly water-efficient (in

fact, vineyards are traditionally considered dryland crops) and second, irrigated vineyards provide quite high economic revenue per hectare.

In the Lower Guadiana basin and TOP domain, on the other hand, irrigated citrus trees and vegetables account for most part of the irrigated area and represent the highest total economic values in this region. What occurs in these two small areas of our study is a general situation in other coastal areas of Andalusia (Hernández-Mora et al. 2001; Vives, 2003).

The economic value of agricultural commodities is an important aspect. For example, many farmers have moved from water-intensive and low economic value crops to water-extensive and higher economic value crops. Alfalfa has been substituted by grapevine or olive trees (Llamas, 2005). According to Llamas (2005) the motto 'more crops and jobs per drop' should be replaced by 'more cash and nature per drop'. Nevertheless, there is still a long way to go to achieve this motto in the Upper and Middle Guadiana basins. In the Lower Guadiana and TOP domain it has been partly achieved, at least on its first half.

Figure 6. Irrigated agriculture green and blue virtual content per crop and year in the different Guadiana sections: UG: Upper Guadiana, MG: Middle Guadiana, LG: Lower Guadiana and TOP domain in different rainfall years (m³/ton). Source: Own elaboration.

D. Agricultural economic productivity (**€**ha)

As it is widely known, agricultural economic productivity of irrigated agriculture is higher than that of rainfed agriculture (Berbel, 2007; Hernández-Mora et al., 2001; MIMAM, 2007). In the case of the Guadiana basin this is true for any type of year (average, humid and dry) (Figure 7). From a socio-economic perspective, irrigated agriculture not only provides a higher income, but also a safer income. This is due both, to the higher

diversification it allows, and to the reduction of climate risks derived from rainfall variability (Comprehensive Assessment of Water Management in Agriculture, 2007).

Concerning the agricultural economic productivity per crop of irrigated agriculture, vegetables have the highest revenues per hectare (5,000-50,000 €ha). Followed by vineyards (about 4,000-6,000 €ha), citrus in the Andalusian section (3,000-5,000 €ha), potatoes (2,000-6,000 €ha) and olive trees (about 1,000-3,000 €ha). Finally grain cereals, grain legumes and industrial crops have productivities of less than 1,000 €ha.

Figure 7. Economic productivity of irrigated and rainfed agriculture per hectare by crop type in the different Guadiana sections in different rainfall years (€ha). Source: Own elaboration.

E. Economic blue water productivity (€m³)

The agricultural total water economic productivity has been calculated in two different ways: using GVA (CHG, 2008b) (Table 1) and using crop economic value (MAPA, 2002) (Figure 8). In both cases the highest value per cubic meter is obtained in the Andalusian part (including the Lower Guadiana and TOP domain), due to the high economic value of the vegetables, which are widespread in the region.

According to Llamas and Martínez-Santos (2005), most probably high value crops are watered with groundwater resources or combining ground and surface water. For instance, Hernández-Mora et al. (2001) show that, in Andalusia (in a study considering almost one million irrigated hectares), agriculture using groundwater is economically over five times more productive and generates almost three times the employment than agriculture using surface water, per unit volume of water used. This difference can be attributed to several causes: the greater control and supply guarantee that groundwater provides, which in turn allows farmers to introduce more efficient irrigation techniques and more profitable crops; the greater dynamism that has characterized the farmer that has sought out his own sources

of water and bears the full costs of drilling, pumping and distribution; and the fact that the higher financial costs farmers bear motivates them to look for more profitable crops that will allow them to maximize their return on investments (Hernández-Mora et al., 2001). Surface and groundwater distinction, therefore, should be taken into account in order to achieve an efficient allocation of water resources. Furthermore, in line with previous studies in arid and semi-arid regions (Garrido et al., 2006; Hernández Mora et al. 2001; Vives 2003), the social (jobs/m³) and economic (€m³) value of groundwater irrigation generally exceeds that of surface water irrigation systems. Agricultural water economic productivity was thus expected to be higher in groundwater based areas.

Along these lines, the Lower Guadiana basin and TOP domain, with a joint surface and groundwater use, have the highest agricultural water economic productivities because they predominantly grow cash crops. The groundwater based Upper Guadiana basin has intermediate values, whereas the surface water based Middle Guadiana shows the lowest water economic productivities. Nevertheless, Upper and Middle Guadiana present similar values in dry years. Probably, this small difference is due on the one hand, to the water irrigation security provided by the existing large surface water reservoirs in the Middle Guadiana; and, on the other, because the use of groundwater in the Upper Guadiana basin has serious legal and political restrictions, at least in theory.

Figure 8. Total blue water economic productivity (€m³) concerning agricultural water consumption by year in the Upper, Middle and Lower Guadiana and TOP domain. Source: Own elaboration.

The water economic productivity analysis can be very useful in order to identify possible water uses not justified in economic efficiency terms and achieve an efficient allocation of water resources.

According to MIMAM (2007), average productivity of blue water used in irrigated agriculture in Spain is about $0.44 \ \text{em}^3$. When looking at the productivity per crop type in the Guadiana basin (Figure 9), vegetables (including horticultural and greenhouse crops) present the highest economic value per water unit (amounting to $15 \ \text{em}^3$ in the Andalusian part: Lower Guadiana and TOP domain). These numbers are similar to the figures estimated by Vives (2003) for greenhouse cultivation using groundwater in Almeria, which amount to $12 \ \text{em}^3$. With lower values vineyards ($1-3 \ \text{em}^3$), potatoes ($0.5-1.5 \ \text{em}^3$), olive tree ($0.5-1 \ \text{em}^3$) and citrus trees ($0.3-0.9 \ \text{em}^3$) show intermediate values. Finally, with remarkably lower values, grain cereals, grain legumes and industrial crops display an average productivity of less than $0.3 \ \text{em}^3$. These data clearly show that the problem in the Guadiana basin is not water scarcity but the use of water for low value crops. Once again, the policy in the near future has to be to more cash per drop.

Figure 9. Blue water economic productivity (€m³) concerning agricultural water consumption by crop and year in the Upper, Middle and Lower Guadiana and TOP domain. Source: Own elaboration.

F. Agricultural trade

The international trade data provided in this section are given at a provincial level as more disaggregated data were not found (ICEX, 2008). The main provinces of each river basin section have been analysed: Ciudad Real for the Upper Guadiana, Badajoz for the Middle Guadiana and Huelva for the Lower Guadiana and TOP domain. Part of the data concerning Ciudad Real and Badajoz were already considered in section 7.1 F.

Concerning trade in tonnes, euros and virtual water, it is noteworthy that Ciudad Real is a net exporter, mainly of wine, and barely imports any commodity (Figure 10). During the

studied period this province has relied on its own food production without depending on global markets. This has been probably at the cost of using its scarce water resources.

In relation to Badajoz, is a net canned-tomato exporter, while importing other commodities such as cereals. It has to be highlighted the increase in cereal imports in drier years (Figure 11).

Huelva also imports virtual water intense commodities, such as cereals, whereas exports low virtual water content fruits (Figure 12). The drier the year the higher the cereal imports. In hydrologic terms, cereal virtual water imports save 1015 Mm³ in Huelva, whereas vegetable exports just uses 100 Mm³. Even if in terms of tonnes and water consumption cereal imports remarkably surpass fruit exports, in economic terms fruit exports are much more important than cereal imports. These results are in line with those obtained by Chahed et al. (2007) when analysing the water footprint in Tunisia, even if they did not assess the economic aspects.

Virtual water imports, and in particular cereal imports, play a role in compensating for the water deficit and providing water and food security in the Middle Guadiana and Andalusian part (Lower Guadiana and TOP domain). For these regions, however, the underlying motivation of importing food (virtual water) is probably hardly a pursuit of comparative advantage, but to fill the domestic shortfall of food supply and to maintain social stability.

According to the World Water Council (2004) one can only speak of virtual water 'trade' if conscious choices are made in water and environmental management policies whether or not to make water available or to release pressure on the domestic water resources by importing goods that else would have consumed much of the domestic water resources available. To make conscious choices, the elements of choice and the players involved in virtual water 'trade' have to be made visible. Allan (2001) states that virtual water 'trade' is

so successful because it is invisible and is applied beyond the general political debate. However, invisibility may lead to postponement of necessary reforms by politicians as imports can be regarded as 'secret reserves' that might bail out in the short run (Warner, 2003).

Finally, the concept of virtual water 'trade' could be very relevant for this region. Local planning and regional collaboration incorporating the notion of virtual water 'trade' could result in exchange of goods, diversification of crops, diet awareness creation or crop replacement actions.

Figure 10. Agricultural commodity export and import in thousand tonnes, million euros and million cubic metres from Ciudad Real during the years 1997 (Humid), 2001 (average) and 2005 (dry). Source: Own elaboration based on ICEX (2008) and Chapagain and Hoekstra (2004) data.

Figure 11. Agricultural commodity export and import in thousand tonnes, million euros and million cubic metres from Badajoz during the years 1997 (Humid), 2001 (average) and 2005 (dry). Source: Own elaboration based on ICEX (2008) and Chapagain and Hoekstra (2004) data.

Figure 12. Agricultural commodity export and import in thousand tonnes, million euros and million cubic metres from Huelva during the years 1997 (Humid), 2001 (average) and 2005 (dry). Source: Own elaboration based on ICEX (2008) and Chapagain and Hoekstra (2004) data.

5.3. Review of crop water consumption estimates by various experts

The present study should be taken as a very interesting but rough approximation to the reality. In tables 2 and 3 green and blue water requirements of the analysed crops by various sources are presented.

When comparing the green water consumption data with other sources, there is a remarkable disparity derived from the methodology in use (Table 2). The present green crop water use numbers, based on FAO Penman-Monteith equation and CROPWAT model,

are higher than figures given by the ITAP (2008), based on the FAO Penman-Monteith equation and an estimation of effective irrigation as 70% of total rainfall. Furthermore, small changes in planting and harvest dates entail big changes in crop water use figures (m³/ha). This could explain these differences.

With regard to the different rainfall years, as expected, there are notable differences depending on the type of year, being lower in dry years (Table 2).

When looking at the theoretical blue water consumption values, the present research results do not seem to differ significantly from other sources (Table 3). As shown in table 2, wheat and other cereals as a whole consume great amounts of blue water whereas their economic value in the markets is very low. Olive tree and vineyard blue water requirements vary depending on the source but they are generally somewhat lower than those of the cereals.

In our opinion, even if these data are a first approximation, they clearly show that the water policy in the Guadiana Basin can and should apply progressively the motto 'more cash and nature per drop'.

6. Conclusions

1. The present virtual water and water footprint analysis, both hydrological and economic, of the Guadiana river basin, provides very interesting results. This analysis however is a first approximation. The calculated theoretical crop water requirements somewhat differ from other authors. There is an outstanding dispersion of data amounting to 100% in certain cases that may be originated by the different methodologies. On the whole, our crop water requirements are based on FAO Penman-Monteith equation and CROPWAT model, whereas figures given by the CHG (2008b) and SIAR (2008) are based on the Thornthwaite and FAO Penman-Monteith equation respectively. In other cases, the uncertainties on some

basic data are related to political issues. One example of this is the lack of acceptable accuracy on the inventory of water users and rights, and on the irrigated area by legal and illegal water wells.

- 2. As in most arid and semiarid regions, in the Guadiana river basin the main green and blue water consuming sector is irrigation, with about 95% of total water consumption in the basin as a whole. Concerning the blue water economic productivity, however, urban water supply and industry values are higher than the corresponding value in agriculture. The multifunctional value of agriculture, however, has to be taken into account. Rainfed agriculture has a high relevance in the Guadiana basin in terms of total hectares. Agricultural economic productivity (ton/ha) and total production (ton/year) of rainfed agriculture, however, are notably lower than that of irrigated agriculture. Thus, even if less in extension, irrigated agriculture produces more tonnes and euros than rainfed agriculture. This economic and social fact explains the political relevance of groundwater irrigation in the Upper Guadiana basin.
- 3. As a whole, high virtual-water low-economic value crops are widespread in the analysed Upper and Middle Guadiana regions. For instance, cereals exhibit virtual water values of 1,000-1,300 m³/ton or even higher in dry years. On the other hand, maize and vegetables (mainly tomato and melons) present the smallest values with around 600 and 100-200 m³/ton respectively, due to their high yields.
- 4. One of the most important contributions of the present report is the analysis of the economic productivity of blue water use for the different crops. In the Upper and Middle Guadiana basin, it seems to range between 0.1-0.2 €m³ for low cost cereals and 1.5-4.5 €m³ for vegetables. These values are relatively small in comparison with the ones obtained in the Andalusian region (Lower Guadiana and TOP domain). In this region, for vegetables

(including horticultural and crops under plastic) using jointly surface and groundwater resources, this value can amount to $15 \notin m^3$. Even with lower figures, vineyards $(1-3 \notin m^3)$ and olive trees $(0.5-1 \notin m^3)$ seem to be profitable crops. As a matter of fact it is widely known that farmers are currently changing their production to vineyards and olive trees. It could be interesting to examine these trends in the near future.

- 5. Nevertheless, we can not fall into the simplification that all the water that is not used for vegetables or trees is wasted water. Factors such as risk diversification, labour or other environmental, social, economic and agronomic reasons have to be taken into account in order to find a balance. The major environmental challenge of agriculture is the preservation of the environment without damaging the agricultural sector economy. The amount of crops and the employment generated in the whole Guadiana basin is producing 'more crops and jobs per drop'. The aim now is to achieve the paradigm 'more cash and nature per drop'. The present results, indicating the low water consumption and high economic value of vegetables, followed by vineyards, is one of the factors that has to be taken into account in order to achieve an efficient allocation of water and economic resources.
- 6. Finally, a first estimation of trade in agricultural products is provided considering the international import-exports at a provincial level. The different sections of the Guadiana basin have different trade strategies. On the one hand, the Upper Guadiana basin is a net exporter, mainly of wine, barely importing any food commodity. On the other, the Lower Guadiana and TOP domain import low-value, high water-consuming cereals, while exporting high-value, low virtual-water content crops such as fruits. This reduces the demand on local (green and blue) water resources that can be used to provide ecological services and other more profitable uses.

Acknowledgements

We wish to thank all the people and institutions that have made this research possible. First, we would like to thank Alberto Garrido, Consuelo Varela, Paula Novo and Roberto Rodriguez. We would also like to thank Professor Arjen Hoekstra for his useful advices. Finally, we can not forget the EU NeWater project and Marcelino Botin Foundation who sponsored this research.

Symbols

Glossary

Blue water – surface and ground water (Hoekstra and Chapagain, 2008).

Blue virtual-water content (V_b) – of a product is the volume of surface or ground water that evaporated as a result of the production of the product. In the case of crop production, the blue water content of a crop is defined as the evaporation of irrigation water from the field. In the cases of industrial production and domestic water supply, the blue water content of the product or service is equal to the part of the water withdrawn from ground or surface water (m^3/ton) (Hoekstra and Chapagain, 2008).

Crop consumptive water use (**CWU**) – is defined as the accumulation of daily evapotranspiration over de complete growing period. It has two components: Green crop water and blue crop consumptive water use (m³/ha) (Hoekstra and Chapagain, 2008).

Crop water requirements (**CWR**) – is defined as the total water needed for evapotranspiration, from planting to harvest for a given crop in a specific climate regime, when adequate soil water is maintained by rainfall and/or irrigation so that it does not limit plant growth and crop yield (mm/time period) (Allen et al., 1998).

Crop water supply – is the quantity of irrigation water, in addition to rainfall, applied to meet a crop's evapotranspiration need and normal crop production. It includes soil evaporation and some unavoidable losses under the given conditions. It is expressed in cubic meters for a crop period (m³/year).

Effective rainfall (P_{eff}) – in irrigation practice, that portion of the total precipitation which is retained by the soil so that it is available for crop production (mm/time period) (FAO, 2008).

External water footprint (WF_e) – is defined as the annual volume of water resources used in other countries or regions to produce goods and services consumed by the inhabitants of

the country or region concerned (km³/year, m³/capita/year) (Hoekstra and Chapagain, 2008).

Green virtual-water content (V_g) – of a product is the volume of rainwater that evaporated during the production process. This is mainly relevant for agricultural products, where it refers to the total rainwater evaporation from the field during the growing period of the crop (including both transpiration by the plants and other forms of evaporation) (m^3 /ton) (Hoekstra and Chapagain, 2008).

Green water – rainwater stored in the soil as soil moisture, also called soil water (Hoekstra and Chapagain, 2008).

Gross value added (**GVA**) – is the value of goods and services produced in an economy at different stages of the productive process (million €). The gross value added is equal to net output or benefit that can be used for the remuneration of productive factors.

Internal water footprint (WF_i) – is defined as the use of domestic water resources to produce goods and services consumed by inhabitants of a country or region (km³/year, m³/capita/year) (Hoekstra and Chapagain, 2008).

Total economic agricultural production – is defined as the total economic value received by the agricultural sector of the region for the commodities sold in the market without taking subsidies into account (total €).

Virtual-water content (V) – the virtual-water content of a product (a commodity, good or service) is the volume of freshwater used to produce the product, measured at the place where the product was actually produced (production-site definition) (m³/ton) (Hoekstra and Chapagain, 2008).

Water footprint (WF) – the water footprint of an individual or community is defined as the total volume of freshwater that is used to produce the goods and services consumed by the individual or community. A water footprint can be calculated for any well-defined group of consumers, including a family, business, village, city, province, state or nation. A water footprint is generally expressed in terms of the volume of water use per year (km³/year, m³/capita/year) (Hoekstra and Chapagain, 2008).

Bibliography

- Aldaya, M.M., Llamas, M. R., Garrido, A. and Varela, C. 2008. Importancia del conocimiento de la huella hidrológica para la política española del agua. *Encuentros Multidisciplinares*, Vol. 10, No. 29, pp. 8-20.
- Aldaya, M.M. and Llamas, M. R. in press. Water footprint analysis for the Guadiana river basin. Value of Water Research Report Series No. 23, UNESCO- IHE. Delft, The Netherlands.
- Allan, J.A. 1997. 'Virtual water': a long term solution for water short Middle Eastern economies? Water Issues Group, School of Oriental and African Studies. University of London. London. [online] Available from:

 http://www.soas.ac.uk/research/our_research/projects/waterissues/papers/occasional-papers.html [Accessed 8 September 2008].
- Allan, J.A. 1999. Water stress and global mitigation: water food and trade. *Aridlands Newsletter*. [online] Available from: http://ag.arizona.edu/OALS/ALN/aln45/allan.html [Accessed 8 September 2008].
- **Allan, J.A. 2001.** Virtual water -economically invisible and politically silent- a way to solve strategic water problems. *International Water and Irrigation*, Vol. 21, No. 4, pp. 39-41.
- Allan, J.A. 2006. Virtual water, part of an invisible synergy that ameliorates water scarcity.

 Water Crisis: Myth or Reality? Ed. Rogers, Llamas and Martinez-Cortina. Balkema
 Publishers.
- Allen, R.G., Pereira, L.S., Raes, D. and Smith, M. 1998. Crop Evapotranspiration Guidelines for Computing Crop Water Requirements FAO Irrigation and drainage paper 56. Food and Agriculture Organization. Rome, Italy.

- Berbel, J. 2007. Análisis Económico del Uso del Agua en la Agricultura y la Ganadería.

 Jornadas de debate sobre El Uso del Agua en la Economía Española. Situación y

 Perspectivas. Grupo de Análisis Económico del Ministerio de Medio Ambiente.

 Sevilla.
- Comprehensive Assessment of Water Management in Agriculture, 2007. Water for Food, Water for Life: A Comprehensive Assessment of Water Management in Agriculture. Earthscan. London.
- **Chahed, J., Besbes, M. and Hamdane, A. 2007.** Stress hydrique et sécurité alimentaire: une vision intégrale des ressources en eau. *La Houille Blanche*, no 3, pp.52-57
- Chapagain, A.K. and Hoekstra, A.Y. 2003. Virtual water flows between nations in relation to trade in livestock and livestock products. Value of Water Research Report Series No. 13, UNESCO- IHE Delft, The Netherlands.
- Chapagain, A.K. and Hoekstra, A.Y. 2004. Water footprints of nations. Value of Water Research Report Series No. 16, UNESCO-IHE, Delft, the Netherlands. [online]

 Available from: http://www.waterfootprint.org/?page=files/Research%20data
 [Accessed 8 September 2008]
- Chapagain, A.K., Hoekstra, A.Y. and Savenije, H.H.G. 2005. Saving water through global trade. Value of Water Research Report Series No. 17, UNESCO- IHE. Delft, The Netherlands.
- Chapagain, A.K., Hoekstra, A.Y., Savenije, H.H.G. and Gautam, R. 2006. The water footprint of cotton consumption: An assessment of the impact of worldwide consumption of cotton products on the water resources in the cotton producing countries. *Ecological Economics*, Vol. 60, No. 1, pp. 186-203.

- Chapagain, A.K. and Orr, S. 2008. An improved water footprint methodology to link global consumption to local water resources: A case study of Spanish tomato consumption. *Journal of Environmental Management*: (In press, doi:10.1016/j.jenvman.2008.1006.1006).
- CHG, 2005. Plan Especial del Alto Guadiana: borrador del documento de directrices.

 Ministerio de Medio Ambiente. Guadiana RBA.
- CHG, 2008a. Confederación Hidrográfica del Guadiana. Guadiana River Basin Authority, Spanish Ministry of Environment [online] Available from: http://www.chguadiana.es/ [Accessed 5 July 2008].
- CHG, 2008b. Información del Análisis Económico de la DHG año 2006, datos de base 2.001. Guadiana River Basin Authority, Spanish Ministry of Environment [online] Available from: http://www.chguadiana.es/ [Accessed 5 July 2008].
- CHG, 2008c. Plan Espacial del Alto Guadiana. Guadiana River Basin Authority, Spanish Ministry of Environment [online] Available from: http://www.chguadiana.es/ [Accessed 5 July 2008].
- **Falkenmark, M. 2003.** Freshwater as shared between society and ecosystems: from divided approaches to integrated challenges. *Philosophical Transactions of the Royal Society B: Biological Sciences*, Vol. 358, No 1440, pp. 2037–2049.
- **Falkenmark, M. and Rockström, J. 2004.** *Balancing Water for Humans and Nature: The New Approach in Ecohydrology*, Earthscan, London, UK.
- FAO, 2003. CROPWAT Model. Food and Agriculture Organization. Rome, Italy.
- **FAO, 2008.** AQUASTAT. *Food and Agriculture Organization*. Rome, Italy. [online]

 Available from: http://www.fao.org/nr/water/aquastat/countries/index.stm [Accessed 8

 September 2008]

- **Garrido A., Martínez-Santos P. and Llamas M.R. 2006.** Groundwater irrigation and its implications for water policy in semiarid countries: the Spanish experience. *Hydrogeology Journal*, Vol. 14, No. 3, pp. 340-349.
- Garrido, A. and Varela-Ortega, C. 2008. Economía del agua en la agricultura e integración de políticas sectoriales. Panel de Estudios. Universidad de Sevilla Ministerio de Medio Ambiente, pp. 1-33.
- Hernández-Mora N., Llamas M.R., Martínez-Cortina L. 2001. Misconceptions in aquifer over-exploitation. Implications for water policy in Southern Europe. Agricultural Use of Groundwater: Towards Integration between Agricultural Policy and Water Resources Management, Dosi C. (ed.), Kluwer Academic Publishers, pp.107–125.
- Hernández-Mora, N., Martinez-Cortina, L. and Fornes, J. 2003. Intensive groundwater use in Spain. M.R. *Intensive Use of Groundwater: Challenges and Opportunities*. Llamas and E. Custodio, eds, Leiden, Netherlands: Balkema.
- Hernández-Mora, N., Martinez-Cortina, L., Llamas, M. R. and Custodio, E. 2007.

 Groundwater Issues in Southern EU Member States. Spain Country Report. [online]

 Available from: http://rac.es/2/2 ficha.asp?id=119&idN3=6&idN4=40 [Accessed 15

 March 2008]
- **Hoekstra, A.Y. 2003.** Virtual water trade between nations: A global mechanism affecting regional water systems. *IGBP Global Change News Letter*, No. 54, pp. 2-4.
- Hoekstra, A.Y. and Chapagain, A.K. 2008. Globalization of water: Sharing the planet's freshwater resources. Blackwell Publishing. Oxford, UK.

- Hoekstra, A.Y. and Hung, P.Q. 2002. Virtual water trade: a quantification of virtual water flows between nations in relation to international crop trade. Value of Water Research Report Series No. 11. UNESCO-IHE. Delft, The Netherlands.
- **Hoekstra, A.Y. and Hung, P.Q. 2005.** Globalisation of water resources: international virtual water flows in relation to crop trade. *Global Environmental Change*. Vol. 15, No. 1, pp. 45-56.
- ICEX, 2008. Spanish Institute for Foreign Trade [online] Available from:

 http://www.icex.es/icex/cda/controller/pageICEX/0,6558,5518394_5518974_5536731

 _0_0_-1,00.html [Accessed 10 June 2008].
- INAG, 2007. Water Institute. Portuguese Ministry of Environment, Spatial Planning and Regional Development [online] Available from: http://www.inag.pt/ [Accessed 8 September 2008].
- INE, 2007. National statistics institute [online] Available from: http://www.ine.es/
 [Accessed 8 September 2008].
- **INM, 2007.** National Institute of Meteorology. Spanish Ministry of Environment [online] Available from: http://www.inm.es/ [Accessed 8 September 2008].
- **ITAP, 2008.** Instituto Técnico Agronómico Provincial. Diputación de Albacete. [online] Available from: http://www.itap.es/ [Accessed 8 September 2008].
- **Llamas, M. R. 2005.** Los colores del agua, el agua virtual y los conflictos hídricos. Discurso inaugural del año 2005-06. *Real Academia de Ciencias Exactas, Físicas y Naturales*. Madrid, pp.30.
- Llamas, R., Fornés, J.M., Hernández-Mora, N. and Martínez Cortina, L. 2001. *Aguas Subterráneas: Retos y Oportunidades*. Mundi-Prensa. Madrid, pp. 529.

- **Llamas, M.R. and Garrido, A. 2007.** Lessons from Intensive Groundwater Use in Spain: Economic and Social Benefits and Conflicts, in the Agricultural Groundwater Revolution: Opportunities and Threats to Development, Giordano and Villholth (eds.), CAB International, Wallingford, U.K., pp. 266-295.
- **Llamas, M.R. and Martínez-Santos, P. 2005.** Intensive groundwater use: Silent revolution and potential source of social conflicts, *Journal of Water Resources Planning and Management, American Society of Civil Engineers*, September-October 2005, pp. 337-341.
- MAPA, 1999. 1T sheets. Spanish Ministry of Agriculture, Fisheries and Food.
- **MAPA, 2001a.** Calendario de siembra, recolección y comercialización, años 1996-1998. Spanish Ministry of Agriculture, Madrid, pp.656.
- MAPA, 2001b. 1T sheets. Spanish Ministry of Agriculture, Fisheries and Food.
- MAPA, 2002. Agro-alimentary Statistics Yearbook. Spanish Ministry of Agriculture, Fisheries and Food [online] Available from:

 http://www.mapa.es/es/estadistica/pags/anuario/Anu_02/indice.asp [Accessed 8 September 2008].
- MAPA ,2006. Agro-alimentary Statistics Yearbook. Spanish Ministry of Agriculture,

 Fisheries and Food [online] Available from:

 http://www.mapa.es/es/estadistica/pags/anuario/Anu_06/indice.asp?parte=2&capitulo=16

 [Accessed 8 September 2008].
- MAPA, 2007. Agro-alimentary Statistics Yearbook. Spanish Ministry of Agriculture,

 Fisheries and Food [online] Available from:

 http://www.mapa.es/es/estadistica/pags/anuario/introduccion.htm [Accessed 8

 September 2008].

- Martínez-Santos, P. 2007. Hacia la gestión adaptable del acuífero de la Mancha Occidental. Doctoral dissertation, University Complutense of Madrid, pp. 334.
- MIMAM, 2007. El agua en la economía española: Situación y perspectivas. Spanish Ministry for the Environment [online] Available from:

 http://circa.europa.eu/Public/irc/env/wfd/library?l=/framework_directive/implementati

 on documents 1/wfd reports/member states/spain/article 5/completo nivel1pdf/ EN

 1.0 &a=d [Accessed 20 May 2008].
- **Rockström, J. 2001.** Green water security for the food makers of tomorrow: windows of opportunity in drought-prone savannahs *Water Science and Technology*, Vol. 43, No. 4, pp. 71-78.
- **Rodríguez, R. 2008.** Cálculo de la huella hidrológica de la agricultura española. MSc thesis. *Polytechnic University of Madrid*, Madrid.
- SIAR 2008. Servicio Integral de Asesoramiento al Regante. Consejería de Agricultura, and Universidad de Castilla La Mancha. [online] Available from: http://crea.uclm.es/siar/index.php [Accessed 20 July 2008].
- Tarjuelo J.M. 2000. Informe sobre el Plan Hidrológico Nacional. Ministerio de Medio Ambiente. Madrid, p. 13 [online] Available from: http://www.mma.es/secciones/agua/pdf/informesphnotr/josem_tarjuelo_martinbenito.p
 df [Accessed 7 January 2008].
- **Vives R. 2003.** Economic and social profitability of water use for irrigation in Andalusia. *Water International*, Vol. 28, No. 3, pp. 326–334.
- Warner, J. 2003. Virtual water virtual benefits. In Hoekstra ed. (2003): Virtual Water Trade; proceedings of the International Expert meeting on Virtual Water Trade; Value of Water Research Report Series no 12; IHE Delft the Netherlands.

World Water Council, 2004. E-Conference Synthesis: Virtual Water Trade - Conscious Choices, March 2004.

Zimmer, D. and Renault, D. 2003. Virtual water in food production and global trade:

Review of methodological issues and preliminary results. Virtual Water Trade:

Proceedings of the International Expert meeting on Virtual Water Trade, Value of Water-Research Rapport Series, no. 12, IHE-Delft, The Netherlands.

Appendixes

Tables

Table 1. Water footprint related to production of the Guadiana basin (year 2001)

TOTAL	. GUADIANA ¹						
	Water footprint	Green	Blue	Total	Per capita	GVA ⁷	Water economic productivity
Population	related to production ⁶	N	/lm³/yea	ır	m³/cap/year	million €	€/m³
1,417,810	Agricultural	2,212	1,827	4,039	2,849	1,096	0.60
	Livestock		22	22	16	286	12.74
	Urban		130	130	91	128 ⁸	0.99 ⁹
	Industrial		20	20	14	1,557	77.90
	Total	2,212	1,999	4,211	2,970	3,068	1.53
UPPER	R GUADIANA ²						
	Water footprint	Green	Blue	Total	Per capita	GVA ⁷	Water economic productivity
Population	related to production ⁶	N	/lm³/yea	ır	m ³ /cap/year	million €	€/m ³
636,721	Agricultural	1,286	928	2,214	3,478	599	0.65
	Livestock		5	5	8	131	25.05
	Urban		55	55	86	54 ⁸	0.99 ⁹
	Industrial		12	12	19	929	77.04
	Total	1,286	1,000	2,286	3,591	1,714	1.71
MIDDI F	E GUADIANA ³						
MIDDEL	Water footprint	Green	Blue	Total	Per capita	GVA ⁷	Water economic productivity
Population	related to production ⁶		lm³/yea		m ³ /cap/year	million €	€/m³
672,534	Agricultural	905	886	1,792	2,664	413	0.47
	Livestock		13	13	20	124	9.30
	Urban		65	65	96	64 ⁸	0.99
	Industrial		6	6	9	485	78.82
	Total	905	970	1,876	2,789	1,086	1.12
TOP⁴							
	Water footprint	Green	Blue	Total	Per capita	GVA ⁷	Water economic productivity
Population	related to production ⁶	M	lm³/yea	r	m ³ /cap/year	million €	€/m³
341,080	Agricultural	74	77	151	444	205	2.66
	Livestock		1	1	3	10	8.57
	Urban		38	38	112	38 ⁸	0.99 ⁹
	Industrial		8	8	24	554	68.62
	Total	74	125	199	583	807	6.47
	CHADIANA ⁵						
LOWER	R GUADIANA ⁵ Water footprint	Green	Rlua	Total	Per canita	GV/Δ^7	Water economic productivity
Population	Water footprint related to		Blue lm³/yea	Total r	Per capita m³/cap/year	GVA ⁷ million €	Water economic productivity €/m³
Population	Water footprint related to production ⁶		lm³/yea	r	m ³ /cap/year	million €	
	Water footprint related to production ⁶ Agricultural	M	lm³/yea 13	r 33	m ³ /cap/year	million €	€/m³ 3.54
Population	Water footprint related to production ⁶ Agricultural Livestock	M	lm³/yea	33 1	m³/cap/year 535 20	million €	€/m³ 3.54 7.42
Population	Water footprint related to production ⁶ Agricultural	M	13 13	r 33	m ³ /cap/year	million € 45 9	€/m³ 3.54

- 1 The Total Guadiana region includes the whole Guadiana river basin excluding the TOP domain. It is not the average of the Upper and Middle Guadiana.
- 2 The Upper Guadiana includes a fraction of Castilla-La Mancha Autonomous region.
- 3 The Middle Guadiana includes a fraction of Extremadura (Badajoz and Cáceres).
- 4 In line with CHG (2008b), TOP region is the Tinto, Odiel and Piedras river basin complementary region.
- 5. TheLower Guadiana region includes the fraction of the basin in Huelva.
- 6 Water footprint related to production by economic sectors.
- 7 Source: CHG (2008b)
- 8 Estimated with data from MIMAM (2007): 0.99 €m³ for urban water supply and sanitation in the Guadiana river basin.
- 9 Source: MIMAM (2007)

Table 2. Green water crop consumptive use values (m³/ha) by different sources.

Upper Guadiana		esent study nd Llamas, 2	2008) ¹	Rodríguez (2008) ²	ITAP (2008) ³	Chapagain and Orr (2008) ⁴
Year	Humid 1997	Average 2001	Dry 2005	2001	2001 (2003) ⁵	Historical time series
Location	Ci	udad Real		Castilla-La Mancha	Albacete	Ciudad Real
Water consumption		CWU _g ⁶		CWU _g ⁶	CWU _g ⁶	CWU _g ⁶
Vineyard	$(1452)^7$	$(854)^7$	$(556)^7$	352	237	
Olive tree	$(1820)^7$	$(1057)^7$	$(664)^7$	665	231	
Oat	1237	1540	318	700		
Wheat	1245	1481	341	867	318	
Barley	1237	1540	318	799	319	
Maize	1254	392	319	594	267	
Tomato	(1156) ⁷	$(298)^7$	(319) ⁷			880

¹ Calculations based on FAO Penman-Monteith equation using CROPWAT model (FAO, 2003) and a time step of 5 days.

² Calculations based on FAO Penman-Monteith equation and a time step of 30 days.

³ Calculations based on FAO Penman-Monteith equation, effective irrigation estimated as 70% of total rainfall.

⁴ Calculations based on FAO Penman-Monteith equation using CROPWAT model (FAO, 2003) and a time step of 5 days.

^{5 2001}data for wheat, barley and maize, while 2003 data for vineyard and olive tree.

⁶ Green consumptive water use (m3/ha)

⁷ Estimated applying a location coefficient for localized irrigation (ET0 * Kc * Kl) following SIAR (2008).

Table 3. Blue water crop consumptive use values (m³/ha) by different sources.

Upper Guadiana		resent stud and Llamas		Rodríguez (2008) ²	CHG (2008b) ³	CHG (2008b) ⁴	CHG (2005) ⁵	Tarjuelo (2000) ⁶	PEAG (CHG, 2008c) ⁷	ITAP (2008) ⁸	SIAR (2008) ⁹	Chapagain and Orr (2008) ¹⁰	Hoekstra and Chapagain (2004) ¹¹
Year	Humid 1997	Average 2001	Dry 2005	2001	2001	2001	2001- 2004	1974-1998	Not specified	2001 (2003) ¹²	2001 (2007) ¹³	Not specified	1997-2001
Location	(Ciudad Rea	I	Castilla-La Mancha	La Mancha	Ciudad Real	Western Mancha	Ciudad Real	Western Mancha	Albacete	La Mancha	Ciudad Real	Spain
Water consumption	CWU _b ¹⁴			CWU _b ¹⁴	CWU _b ¹⁴	CWU _b ¹⁴	CWU _b ¹⁴	CWU _b ¹⁴	CWU _b ¹⁴	CWU _b ¹⁴	CWU _b ¹⁴	CWU _b ¹⁴	CWU ¹⁵
Vineyard	(1670) ¹⁶	(2890) ¹⁶	(3619) ¹⁶	3977	2690	3678	1516	2000-2500	3678	2388	1693		6622
Olive tree	(1186) ¹⁶	$(2502)^{16}$	(3271) ¹⁶	3991	1930		2153			2186			7350
Oat	2079	2200	3743	3801	$(2350)^{17}$	2306			2306				2830
Wheat	2058	2277	3759	2533		2583	3342	2842	2583	3902	2403		3070
Barley	2079	2200	3743	3976		2999	2690	2759	2999	2630	1880		2831
Maize	4445	6534	7460	7347		7014	8117	5174	7014	7262	7604		6116
Tomato	(3845) ¹⁶	(5779) ¹⁶	(6510) ¹⁶		(3510) ¹⁸						5705	3730	3165

¹ Calculations based on FAO Penman-Monteith equation using CROPWAT model (FAO, 2003) and a time step of 5 days.

² Calculations based on FAO Penman-Monteith equation and a time step of 30 days.

³ Calculations based on Thornthwaite method.

⁴ Calculations based on FAO Penman-Monteith equation. Do not consider deficit irrigation strategies. These data may vary with respect to other CHG data calculated according to Thornwaite method.

⁵ Calculations following SIAR.

⁶ Calculations based on FAO Penman-Monteith equation, and 25 year climate series. For the vineyard deficit irrigation recommendations are followed (riego deficitario controlado, RDC).

⁷ Source: Tragsatec and MIMAM. Do not consider deficit irrigation strategies.

⁸ Calculations based on FAO Penman-Monteith equation, effective irrigation estimated as 70% of total rainfall.

⁹ Calculations based on FAO Penman-Monteith equation using deficit irritation for trees.

¹⁰ Calculations based on FAO Penman-Monteith equation using CROPWAT model (FAO, 2003) and a time step of 5 days.

¹¹ Calculations based on FAO Penman-Monteith equation using CROPWAT model (FAO, 2003).

^{12 2001}data for wheat, barley and maize, while 2003 data for vineyard and olive tree.

^{13 2001} data for every crop except for tomato (industry) in 2007.

¹⁴ Blue consumptive water use (m³/ha)

¹⁵ Total consumptive water use (including green and blue) (m³/ha)

¹⁶ Estimated applying a location coefficient for localized irrigation (ET₀ * K_c * K_l) following SIAR (2008).

¹⁷ Value for grain cereals

¹⁸ Value for vegetables

Symbols

Symbol	Unit	Description
CWR _[c]	m ³ /year	Crop water requirement of crop c
$\mathrm{CWC}_{[\mathrm{c}]}$	m ³ /ha/year	Crop water consumption to produce a particular crop c, also called evapotranspirative demand
ET_0	mm/day	Reference evapotranspiration
ETc	mm/day	Crop evapotranspiration of a crop c
GVA	million €	Gross Value Added
Kc		Crop coefficient
V	m ³ /ton	Virtual Water Content
V_b	m ³ /ton	Blue Virtual Water Content
$ m V_{g}$	m ³ /ton	Green Virtual Water Content
WF	m ³ /year	Water Footprint
WF_b	m ³ /year	Blue Water Footprint
WF_g	m ³ /year	Green Water Footprint
WF_i	m ³ /year	Internal Water Footprint

Appendixes

A. Crop Area, Production and Yield (2001)

B) UPPER GUADIANA ¹	Α	rea (ha) ²		Prod	uction (ton/	year) ³	Yield	(kg/ha) ⁴
Crops	Rainfed	Irrigated	Total	Rainfed	Irrigated	Total	Rainfed	Irrigated
Grain cereals ⁵ :	478.572	97.634	576.206	1.959.752	740.309	2.700.061	4095	7583
Cereal - Wheat, barley, oat							1045	3460
Cereal – Maize							7145	11705
Grain legumes - Veza, yeros	68.974	10.567	79.541	17.864	11.835	29.699	259	1120
Potatoes	411	733	1.143	4.986	17.855	22.842	12140	24369
Industrial crops – Sunflower	73.038	10.450	83.488	24.541	16.752	41.293	336	1603
Fodder – Veza, alfalfa	30.312	7.701	38.013	182.784	173.269	356.052	6030	22500
Vegetables – melon	488	13.337	13.826	3.959	369.447	373.406	8110	27700
Flowers and ornamental plants	133	100	234					
Seeds and small plants	0	21	21					
Other grass crops	59	1.895	1.954					
Fallow land	343.142	0	343.142					
Vegetable gardens	0	39	39					
Citrus	0	10	10				-	-
Temperate climate fruit trees	84	210	295					
Subtropical climate fruit trees	0	0	0					
Dry fruit trees	5.503	293	5.796					
Olive tree – for olive oil	134.687	13.213	147.900	234.086	31.116	265.202	1738	2355
Vineyard – for wine production	199.277	131.866	331.143	799.100	1.588.985	2.388.085	4010	12050
Nursery	0	25	25					
Other permanent crops	185	6	191					
Greenhouse tree crops	0	2	2					
Mushrooms		15	15					
Greenhouses		86	86					
Total	1.334.865	288.205	1.623.070	3.227.072	2.949.568	6.176.640	4590	12410
	Surface ⁶	26.390						

Surface° 26.390 groundwater ⁷ 237.857

¹ The Upper Guadiana includes a fraction of Castilla-La Mancha Autonomous region.

² Source: CHG (2008b)

³ Calculated multiplying area (CHG, 2008b) and yield (MAPA, 2002)

⁴ *Source*: MAPA (2007)

⁵ Grain cereals comprise main cereals in the area according to MAPA 1T sheets.

⁶ Irrigated area with surface water

⁷ Irrigated area with groundwater

C) MIDDLE GUADIANA ¹	Are	ea (ha) ²		Proc	duction (ton/	year) ³	Yield	(kg/ha) ⁴
Crops	Rainfed	Irrigated	Total	Rainfed	Irrigated	Total	Rainfed	Irrigated
Grain cereals ⁵ :	281.182	96.161	377.343	750.101	742.533	1.492.634	2668	7722
Cereal - Wheat, barley, oat							2668	3734
Cereal – Maize							-	12000
Cereal – Rice							-	7431
Grain legumes – Chick peas	19.535	1.532	21.067	14.651		14.651	750	-
Potatoes	6	430	436	-	10.758	10.758	-	25000
Industrial crops - Sunflower	16.372	15.541	31.913	14.964	44.510	59.474	914	2864
Fodder – Clover, veza	46.957	2.558	49.514	469.565	0	469.565	10000	-
Vegetables – Tomato	504	21.597	22.101	0	1.232.619	1.232.619	-	57073
Flowers and ornamental plants	6	62	68					
Seeds and small plants	0	77	77					
Other grass crops	2	1.859	1.861					
Fallow land	143.481	0	143.481					
Vegetable gardens	0	88	88					
Citrus	0	41	41	0	489	489	-	12000
Temperate climate fruit trees	5.060	7.807	12.867					
Subtropical climate fruit trees	0	3	3					
Dry fruit trees	1.716	1.005	2.721					
Olive tree for olive oil and table	190.661	47.778	238.439	328.700	238.891	567.591	1724	5000
Vineyard for wine production	59.116	11.704	70.819	299.362	93.630	392.992	5064	8000
Nursery	0	69	69					
Other permanent crops	67	0	67					
Greenhouse tree crops	0	4	4					
Mushrooms		0	0					
Greenhouses		77	77					
Total	764.664	208.393	973.057	1.877.343	2.363.430	4.240.774	2950	14082
	Surface ⁶	121.291						

1 The Middle Guadiana includes a fraction of Extremadura (Badajoz and Cáceres).

23.061

2 *Source*: CHG (2008b)

3 Calculated multiplying area (CHG, 2008b) and yield (MAPA, 2002)

groundwater 7

4 *Source*: MAPA (2007)

5 Grain cereals comprise main cereals in the area according to MAPA 1T sheets.

6 Irrigated area with surface water

7 Irrigated area with groundwater

D) TOP ¹	Are	a (ha) ²		Produ	uction (ton	/year) ³	Yield	(kg/ha) ⁴
Crops	Rainfed	Irrigated	Total	Rainfed	Irrigated	Total	Rainfed	Irrigated
Grain cereals ⁵ – Wheat	23,771	1,221	24,992	58,002	4,188	62,190	2440	3430
Grain legumes – Lupin, veza	477	206	683	324	299	623	679	1450
Potatoes	41	121	162	353	1,667	2,020	8554	13807
Industrial crops – Sunflower	10,659	2,579	13,237	11,192	5,286	16,478	1050	2050
Fodder – Veza	809	234	1,043	12,948		12,948	16000	-
Vegetables – Strawberry	131	4,374	4,505	0	147,600	147,600	0	33741
Flowers and ornamental plants	0	66	66					
Seeds and small plants	0	1	1					
Other grass crops	0	0	0					
Fallow land	18,900	0	18,900					
Vegetable gardens	0	27	27					
Citrus	0	7,665	7,665		118,337	118,337	-	15,438
Temperate climate fruit trees	292	1,789	2,081					
Subtropical climate fruit trees	0	101	101					
Dry fruit trees	1,787	81	1,868					
Olive tree for olive oil and table	10,171	1,059	11,229	8,747	1,673	10,420	860	1,580
Vineyard for wine and grape	3,178	129	3,307	23,549	1,056	24,605	7,410	8,200
Nursery	0	6	6					
Other permanent crops	2	0	2					
Greenhouse tree crops	0	64	64					
Mushrooms		0	0					
Greenhouses		352	352					
Total	70,220	20,073	90,293	115,115	280,106	395,221	5285	9962
	Surface ⁶	11.076		· · · · · · · · · · · · · · · · · · ·		•	•	•

1 In line with CHG (2008b), TOP region is the Tinto, Odiel and Piedras river basin complementary region.

8,695

2 *Source*: CHG (2008b)

3 Calculated multiplying area (CHG, 2008b) and yield (MAPA, 2002)

groundwater 7

4 *Source*: MAPA (2007)

5 Grain cereals comprise main cereals in the area according to MAPA 1T sheets.

6 Irrigated area with surface water

7 Irrigated area with groundwater

D) LOWER GUADIANA ¹	Are	ea (ha) ²		Produ	uction (ton	/year)³	Yield	(kg/ha) ⁴
Crops	Rainfed	Irrigated	Total	Rainfed	Irrigated	Total	Rainfed	Irrigated
Grain cereals ⁵ – Wheat	7,363	203	7,566	17,965	696	18,661	2,440	3,430
Grain legumes – Lupin, veza	121	11	132	82	16	98	679	1,450
Potatoes	17	5	22	147	64	210	8,554	13,807
Industrial crops – Sunflower	340	127	468	358	261	618	1,050	2,050
Fodder – Veza	779	234	1,012	12,457		12,457	16,000	-
Vegetables – Strawberry	23	380	403	0	12,817	12,817	-	33,741
Flowers and ornamental plants	0	7	7					
Seeds and small plants	0	0	0					
Other grass crops	0	0	1					
Fallow land	10,839	0	10,839					
Vegetable gardens	0	20	20					
Citrus	0	1,672	1,672		25,817	25,817	-	15,438
Temperate climate fruit trees	104	360	464					
Subtropical climate fruit trees	0	1	1					
Dry fruit trees	3,433	24	3,456					
Olive tree for olive oil and table	5,324	246	5,570	4,579	388	4,967	860	1,580
Vineyard for wine and grape	63	251	314	465	2,061	2,526	7,410	8,200
Nursery	0	0	0					
Other permanent crops	0	0	0					
Greenhouse tree crops	0	0	0					
Mushrooms		0	0					
Greenhouses		7	7					
Total	28,406	3,548	31,954	36,053	42,119	78,171	5,285	9,962
	Surface6	2,435						

1 The Lower Guadiana basin comprises the Guadiana basin part in Huelva.

2 Source: CHG (2008b)

3 Calculated multiplying area (CHG, 2008b) and yield (MAPA, 2002)

groundwater 7

4 *Source*: MAPA (2007)

5 Grain cereals comprise main cereals in the area according to MAPA 1T sheets.

780

6 Irrigated area with surface water

7 Irrigated area with groundwater

B. Crop water consumption (m³/ha), total water resource consumption (10^6 m³/year) and virtual water content (m³/ton) (2001).

	Crop	Water C	onsump	tion (m ³ /	ha)	Total u	se of wa	ter resour	ces (10 ⁶ r	n³/year)	Virtua	l Wate	r Conten	t (V) (m ³	/ton)
B) UPPER GUADIANA ¹	Rainfed		Irrig	ated		Rainfed		Irrig	ated		Rainfed		Irriç	gated	
Crop	V_g^2	V_g^2	V _b ³	V_b^{4}	Total	$V_g^{2*}A^5$	$V_g^{2*}A^5$	$V_b^{3*}A^5$	$V_b^{4*}A^5$	Total*A ⁵	V_g^2	V _g ²	V_b^3	V_b^4	Total
Grain cereals ⁶ :	1238	1238	3303	2599	4541	593	121	322	254	443	302	163	436	343	599
Cereal - Wheat, barley, oat	1520	1520	2225		3746						1455	439	643		1083
Cereal - Maize	392	392	6534		6926						55	34	558		592
Grain legumes Veza, yeros	911	911	2598	2254	3510	63	10	27	24	37	3519	814	2320	2012	3133
Potatoes	370	370	6035	2864	6404			4	2	5	30	15	248	118	263
Industrial crops – Sunflower	311	311	5625	3168	5936	23	3	59	0	62	924	194	3509	0	3703
Fodder – Veza, alfalfa	816	816	4177	4079	4993	25	6	32	31	38	135	36	186	181	222
Vegetables – Melon	290	290	5136	3741	5426	0	4	69	50	72	36	10	185	135	196
Flowers, ornamental plants				4052					0						
Seeds and small plants				3400					0						
Other grass crops				3880					7						
Fallow land									0						
Vegetable gardens				3906					0						
Citrus				3900					0						
Temperate climate fruit trees				3980					1						
Subtropical climate fruit trees															
Dry fruit trees				4915					1						
Olive tree for olive oil	1057	1057	2502	1893	3560	142	14	33	25	47	608	449	1063	804	1512
Vineyard for wine production	854	854	2890	2692	3744	170	113	381	355	494	213	71	240	223	311
Nursery				3400											
Other permanent crops				4047											
Greenhouse tree crops				3400											
Mushrooms				18000											
Greenhouses ⁶				4200											
Total	731	731	4033	2932	4764	1016	271	928	752	1199	728	223	939	477	1161

¹ The Upper Guadiana includes a fraction of Castilla-La Mancha Autonomous region.

² V_g : Green virtual water. *Source*: Own elaboration. 3 V_b : Blue virtual water. *Source*: Own elaboration.

⁴ V_b: Blue virtual water. Source: CHG (2008b)

⁵ A: Area in hectares

⁶ Grain cereals comprise main cereals in the area according to MAPA 1T sheets.

	Crop Water Consumption (m³/ha)					Total us	se of wat	er resou	rces (10	⁵ m ³ /year)	Virtual Water Content (V) (m³/ton)				
C) MIDDLE GUADIANA ¹	Rainfed		Irriga	ited		Rainfed		Irri	gated		Rainfed		Irrig	ated	
Crop	V_g^2	V_g^2	V _b ³	V _b ⁴	Total	$V_g^{2*}A^5$	$V_g^{2*}A^5$	$V_b^{3*}A^5$	$V_b^{4*}A^5$	Total*A ⁵	V_g^2	V _g ²	V _b ³	V_b^4	Total
Grain cereals ⁶ :	1378	1052	4462	4.095	5514	387	101	429	394	530	516	136	578	530	714
Cereal - Wheat, barley, oat	1378	1378	2473		3851						516	369	662		1031
Cereal - Maize	-	366	6712		7078							30	559		590
Cereal - Rice	-	760	8178		8938							102	1100		1203
Grain legumes – Chick peas	325	-	-	3.050		6	0	0	5	0	433				
Potatoes	-	970	3437	2.821	4406	0	0	1	1	2					
Industrial crops – Sunflower	325	325	5741		6065	5	5	89	0	94	355	113	2004	0	2118
Fodder - Clover, veza	1665	1665	1745	5.346		78			14	0	167				
Vegetables – Tomato	317	317	6592	4.043	6909	0	7	142	87	149	-	6	115	71	121
Flowers, ornamental plants				4.050					0						
Seeds and small plants				3.400					0						
Other grass crops				4.430					8						
Fallow land															
Vegetable gardens				3.637					0						
Citrus	-	2244	6000	3.900	8244		0	0	0			187	500	325	687
Temperate climate fruit trees				3.718					29						
Subtropical climate fruit trees				4.000					0						
Dry fruit trees				5.500					6						
Olive tree	1048	1048	3733	1.975	4781	200	50	178	94	228	608	210	747	395	956
Vineyard	912	912	3901	2.683	4814	54	11	46	31	56	180	114	488	335	602
Nursery				3.400											
Other permanent crops															
Greenhouse tree crops				3.400											
Mushrooms															
Greenhouses				4.200											
Total	853	1067	4451	3758	5819	731	174	886	671	1061	397	141	750	276	891

- 1 The Middle Guadiana includes a fraction of Extremadura (Badajoz and Cáceres).
- 2 V_g: Green virtual water. *Source*: Own elaboration. 3 V_b: Blue virtual water. *Source*: Own elaboration. 4 V_b: Blue virtual water. *Source*: CHG (2008b)

- 5 A: Area in hectares
- 6 Grain cereals comprise main cereals in the area according to MAPA 1T sheets.

	Crop Water Consumption (m³/ha) Rainfed Irrigated					Total u	se of wa	er resourc	ces (10 ⁶ r	n³/year)	Virtual	l Wate	r Content	(V) (m ³	/ton)
D) TOP ¹	Rainfed		Irriga	ted		Rainfed		Irrig	ated		Rainfed		Irriga	ted	
Crop	V_g^2	V _g ²	V_b^3	V_b^4	Total	$V_g^{2*}A^5$	$V_g^{2*}A^5$	$V_b^{3*}A^5$	$V_b^{4*}A^5$	Total*A ⁵	V_g^2	V_g^2	V_b^3	V_b^4	Total
Grain cereals ⁶ – Wheat	1380	1380	2760	3677	4140	33	2	3	4	5	565	402	805	1072	1207
Grain legumes - Lupin, veza	752	752	4227	3050	4979	0	0	1	1	1	1108	519	2915	2103	3434
Potatoes	1015	1015	3560	1240	4575	0	0	0	0	1	119	74	258	90	331
Industrial crops – Sunflower	0	0	5936		5936	0	0	15	0	15	0	0	2896	0	2896
Fodder – Veza and others	1505	1505	3674	5023	5178	1			1		94				
Vegetables – Strawberry	1688	1688	2836	3840	4523	0	7	12	17	20	-	50	84	114	134
Flowers and ornamental plants				4050					0						
Seeds and small plants				3400					0						
Other grass crops															
Fallow land															
Vegetable gardens				3817					0						
Citrus	-	2828	5586	3952	8415		22	43	30	65		183	362	256	545
Temperate climate fruit trees				3765					7						
Subtropical climate fruit trees				4000					0						
Dry fruit trees				5900					0						
Olive tree for oil and table	589	589	1601	2282	2189	6	1	2	2	2	685	373	1013	1444	1386
Vineyard for wine and grape	564	564	1902	2888	2466	2	0	0	0	0	76	69	232	352	301
Nursery				3400					0						
Other permanent crops															
Greenhouse tree crops				3400					0						
Mushrooms															
Greenhouses ⁶				4200					1_						
Total	936	1147	3565	3635	4711	42	32	77	66	109	378	209	1071	679	1279

- 1 In line with CHG (2008b), TOP region is the Tinto, Odiel and Piedras river basin complementary region.
- 2 V_g: Green virtual water. *Source*: Own elaboration.
 3 V_b: Blue virtual water. *Source*: Own elaboration.
 4 V_b: Blue virtual water. *Source*: CHG (2008b)

- 5 A: Area in hectares
- 6 Grain cereals comprise main cereals in the area according to MAPA 1T sheets.

	Crop Water Consumption (m³/ha) Rainfed Irrigated					Total u	se of wa	er resourc	ces (10 ⁶ r	n³/year)	Virtual	Wate	r Content	(V) (m ³	/ton)
D) LOWER GUADIANA ¹	Rainfed		Irriga	ted		Rainfed		Irrig	ated		Rainfed		Irriga	ted	
Crop	V_g^2	V_g^2	V_b^3	V_b^4	Total	$V_g^{2*}A^5$	$V_g^{2*}A^5$	$V_b^{3*}A^5$	V _b ⁴ *A ⁵	Total*A ⁵	V_g^2	V_g^2	V_b^3	V_b^4	Total
Grain cereals ⁶ – Wheat	1380	1380	2760	3677	4140	10	0	1	1	1	565	402	805	1072	1207
Grain legumes - Lupin, veza	752	752	4227	3050	4979	0	0	0	0	0	1108	519	2915	2103	3434
Potatoes	1015	1015	3560	1240	4575	0	0	0	0	0	119	74	258	90	331
Industrial crops – Sunflower	0	0	5936		5936	0	0	1	0	1	0	0	2896	0	2896
Fodder – Veza and others	1505	1505	3674	5023	5178	1			1		94				
Vegetables – Strawberry	1688	1688	2836	3840	4523	0			1	2	-	50	84	114	134
Flowers and ornamental plants				4050					0						
Seeds and small plants				3400					0						
Other grass crops															
Fallow land															
Vegetable gardens				3817					0						
Citrus	-	2828	5586	3952	8415	0	5	9	7	14		183	362	256	545
Temperate climate fruit trees				3765					1						
Subtropical climate fruit trees				4000					0						
Dry fruit trees				5900					0						
Olive tree for oil and table	589	589	1601	2282	2189	3	0	0	1	1	685	373	1013	1444	1386
Vineyard for wine and grape	564	564	1902	2888	2466	0	0	0	1	1	76	69	232	352	301
Nursery				3400					0						
Other permanent crops															
Greenhouse tree crops				3400					0						
Mushrooms															
Greenhouses ⁶				4200					0						
Total	936	1147	3565	3635	4711	15	6	13	13	19	378	209	1071	679	1279

The Lower Guadiana basin comprises the Guadiana basin part in Huelva.
 V_g: Green virtual water. *Source*: Own elaboration.
 V_b: Blue virtual water. *Source*: Own elaboration.
 V_b: Blue virtual water. *Source*: CHG (2008b)

⁵ A: Area in hectares

⁶ Grain cereals comprise main cereals in the area according to MAPA 1T sheets.

C. Economic value and economic water productivity (€m³) (2001)

	Economic value				Water economic productivity ⁵	GVA ⁶	Employment ⁷	
B) UPPER GUADIANA ¹	€/ha²	€/ha²	€/ton³	Total million € ⁴		€/m³	million €	post number
Crop	Rainfed	Irrigated		Rainfed	Irrigated	Irrigated		
Grain cereals ⁸ :	549	1017	134	263	99	0,3		
Cereal - Wheat, barley, oat			133					
Cereal - Maize			136					
Grain legumes – Veza, yeros	46	197	176	3	2	0,1		
Potatoes	2508	5035	207	1	4	0,8		
Industrial crops – Sunflower		410	256		4	0,1		
Fodder – Veza, alfalfa			101					
Vegetables – Melon	2092	7144	258	1	95	1,4		
Flowers, ornamental plants								
Seeds and small plants								
Other grass crops								
Fallow land								
Vegetable gardens								
Citrus			192					
Temperate climate fruit trees								
Subtropical climate fruit trees								
Dry fruit trees								
Olive tree for olive oil	865	1172	498	116	15	0,5		
Vineyard for wine production	1823	5479	455	363	722	1,9		
Nursery								
Other permanent crops								
Greenhouse tree crops								
Mushrooms								
Greenhouses								
Total	560	3271		748	943	1,0	599	26818

- 1 The Upper Guadiana includes a fraction of Castilla-La Mancha Autonomous region.
- 2 Total economic value (total €) divided by area (ha)
- 3 Source: MAPA (2002)
- 4 Economic value (€ton) multiplied by production (ton/year). *Source*: MAPA (2002)
- 5 Total economic value (total €) divided by the total use of water resources (m3/year)
- 6 Gross Value Added (GVA). Source: CHG (2008b)
- 7 *Source*: CHG (2008b)
- 8 Grain cereals comprise main cereals in the area according to MAPA 1T sheets.

	Economic value				Water economic productivity ⁵	GVA ⁶	Employment ⁷	
C) MIDDLE GUADIANA1	€/ha²	€/ha²	€/ton ³	Total million €⁴		€/m³	million €	post number
Crop	Rainfed	Irrigated		Rainfed	Irrigated	Irrigated		
Grain cereals8:	435	1259	163	122	121	0,3		
Cereal - Wheat, barley, oat			133					
Cereal - Maize			136					
Cereal - Rice			279					
Grain legumes – Chick peas	613		817	12				
Potatoes		5165	207		2	1,5		
Industrial crops – Sunflower		732	256		11	0,1		
Fodder - Clover, veza	0		101					
Vegetables – Tomato	0	19182	336	0	414	2,9		
Flowers, ornamental plants								
Seeds and small plants								
Other grass crops								
Fallow land								
Vegetable gardens								
Citrus		2302	192		0	0,4		
Temperate climate fruit trees								
Subtropical climate fruit trees								
Dry fruit trees								
Olive tree for oil and table	858	2488	498	164	119	0,7		
Vineyard for wine production	2303	3638	455	136	43	0,9		
Nursery								
Other permanent crops								
Greenhouse tree crops								
Mushrooms								
Greenhouses								
Total	568	3409		434	711	0,8	413	22991

- 1 The Middle Guadiana includes a fraction of Extremadura (Badajoz and Cáceres).
- 2 Total economic value (total €) divided by area (ha)
- 3 *Source*: MAPA (2002)
- 4 Economic value (€ton) multiplied by production (ton/year). *Source*: MAPA (2002) 5 Total economic value (total €) divided by the total use of water resources (m3/year)
- 6 Gross Value Added (GVA). Source: CHG (2008b)
- 7 *Source*: CHG (2008b)
- 8 Grain cereals comprise main cereals in the area according to MAPA 1T sheets.

	Economic value				Water economic productivity⁵	GVA ⁶	Employment ⁷	
D) TOP ¹	€/ha²	€/ha²	€/ton ³	Total million € ⁴		€/m³	million €	post number
Crop	Rainfed	Irrigated		Rainfed	Irrigated	Irrigated		
Grain cereals ⁸ – Wheat	325	457	133	8	1	0,2		
Grain legumes – Lupin, veza	129	276	190	0	0	0,1		
Potatoes	1767	2852	207	0	0	0,8		
Industrial crops – Sunflower		524	256		1	0,1		
Fodder – Veza and others			101					
Vegetables - Strawberries	0	28039	831	0	123	9,9		
Flowers and ornamental plants								
Seeds and small plants								
Other grass crops								
Fallow land								
Vegetable gardens								
Citrus		2961	192		23	0,5		
Temperate climate fruit trees								
Subtropical climate fruit trees								
Dry fruit trees								
Olive tree	428	786	498	4	1	0,5		
Vineyard	3369	3728	455	11	0	2,0		
Nursery								
Other permanent crops								
Greenhouse tree crops								
Mushrooms								
Greenhouses ⁶								
Total	327	7422		23	149	1,9	205	9435

- 1 In line with CHG (2008b), TOP region is the Tinto, Odiel and Piedras river basin complementary region.
- 2 Total economic value (total €) divided by area (ha)
- 3 *Source*: MAPA (2002)
- 4 Economic value (€ton) multiplied by production (ton/year). *Source*: MAPA (2002)
- 5 Total economic value (total €) divided by the total use of water resources (m3/year)
- 6 Gross Value Added (GVA). Source: CHG (2008b)
- 7 *Source*: CHG (2008b)
- 8 Grain cereals comprise main cereals in the area according to MAPA 1T sheets.

	Economic value				Water economic productivity⁵	GVA ⁶	Employment ⁷	
D) LOWER GUADIANA ¹	€/ha²	€/ha²	€/ton ³	Total million € ⁴		€/m³	million €	post number
Crop	Rainfed	Irrigated		Rainfed	Irrigated	Irrigated		
Grain cereals ⁸ – Wheat	325	457	133	2	0	0,2		
Grain legumes – Lupin, veza	129	276	190	0	0	0,1		
Potatoes	1767	2852	207	0	0	0,8		
Industrial crops – Sunflower		524	256		0	0,1		
Fodder – Veza and others			101					
Vegetables - Strawberries	0	28039	831	0	11	9,9		
Flowers and ornamental plants								
Seeds and small plants								
Other grass crops								
Fallow land								
Vegetable gardens								
Citrus		2961	192		5	0,5		
Temperate climate fruit trees								
Subtropical climate fruit trees								
Dry fruit trees								
Olive tree	428	786	498	2	0	0,5		
Vineyard	3369	3728	455	0	1	2,0		
Nursery								
Other permanent crops								
Greenhouse tree crops								
Mushrooms								
Greenhouses ⁶								
Total	174	4765		5	17	1,3	45	2206

- 1 The Lower Guadiana basin comprises the Guadiana basin part in Huelva.
- 2 Total economic value (total €) divided by area (ha)
- 3 *Source*: MAPA (2002)
- 4 Economic value (€ton) multiplied by production (ton/year). *Source*: MAPA (2002)
- 5 Total economic value (total €) divided by the total use of water resources (m3/year)
- 6 Gross Value Added (GVA). Source: CHG (2008b)
- 7 Source: CHG (2008b)
- 8 Grain cereals comprise main cereals in the area according to MAPA 1T sheets.

Figures




Figure 1. Guadiana river basin geographic and administrative domain from 1 January 2006 onwards (CHG, 2008a)




Figure 2. Western Mancha aquifer location within the Upper Guadiana Basin. Modified from CHG (2008b).




Figure 3. Percentage of areas of irrigated and rainfed crops in the Upper, Middle, Lower Guadiana and TOP domain (average-year 2001). Showing crops occupying over 1% of land. *Source*: CHG (2008b).




Figure 4. Theoretical green and blue agricultural water consumption (Mm³/year) in the Upper, Middle, Lower Guadiana and TOP domain a dry (2005), average (2001) and humid year (1997). *Source*: Own elaboration.




Figure 5. Theoretical green and blue (surface and ground) agricultural water consumption (Mm³/year) in the Upper, Middle, Lower Guadiana and TOP domain a dry (2005), average (2001) and humid year (1997). The size of the circle is proportional to the volume of water. *Source*: Own elaboration.




Figure 6. Irrigated agriculture green and blue virtual content per crop and year in the different Guadiana sections: UG: Upper Guadiana, MG: Middle Guadiana, LG: Lower Guadiana and TOP domain in different rainfall years (m³/ton). *Source*: Own elaboration.




Figure 7. Economic productivity of irrigated and rainfed agriculture per hectare by crop type in the different Guadiana sections in different rainfall years (€ha). *Source*: Own elaboration.

Water economic productivity




Figure 8. Total blue water economic productivity $(\not\in m^3)$ concerning agricultural water consumption by year in the Upper, Middle and Lower Guadiana and TOP domain. *Source*: Own elaboration.




Figure 9. Blue water economic productivity $(\not\in m^3)$ concerning agricultural water consumption by crop and year in the Upper, Middle and Lower Guadiana and TOP domain. *Source*: Own elaboration.




Figure 10. Agricultural commodity export and import in thousand tonnes, million euros and million cubic metres from Ciudad Real during the years 1997 (Humid), 2001 (average) and 2005 (dry). *Source*: Own elaboration based on ICEX (2008) and Chapagain and Hoekstra (2004) data.




Figure 11. Agricultural commodity export and import in thousand tonnes, million euros and million cubic metres from Badajoz during the years 1997 (Humid), 2001 (average) and 2005 (dry). *Source*: Own elaboration based on ICEX (2008) and Chapagain and Hoekstra (2004) data.




Figure 12. Agricultural commodity export and import in thousand tonnes, million euros and million cubic metres from Huelva during the years 1997 (Humid), 2001 (average) and 2005 (dry). *Source*: Own elaboration based on ICEX (2008) and Chapagain and Hoekstra (2004) data.